

**11th International Meeting on Statistical Climatology
Session 10: Long Term Memory**

University of Edinburgh, Scotland, 12-16 July, 2010

Memory in climate and things not to be forgotten



Demetris Koutsoyiannis

Department of Water Resources and Environmental Engineering
Faculty of Civil Engineering
National Technical University of Athens, Greece
(dk@itia.ntua.gr, <http://www.itia.ntua.gr/dk/>)



Presentation available online: <http://www.itia.ntua.gr/en/docinfo/991/>

1. What is memory, short memory and long memory?

What do books say about “memory”?

- Papoulis (1991; Probability, Random Variables, and Stochastic Processes \equiv my gospel) does not use the term **memory** for a stochastic process
 - He only defines **memoryless systems**, as those whose outputs $y(t)$ are given as a function $g(x(t))$ of the input $x(t)$ at the same time t
 - This involves two stochastic processes and cannot be applied to a single stochastic process
- Beran (1994, Statistics for Long-Memory Processes) defines long and short memory **intuitively** and relates it to autocorrelation $\rho(j)$ for lag j :
 - “The intuitive interpretation of $[\rho(j) \approx c_\rho |j|^{-\alpha}]$ is that the process has **long memory**”
 - “in contrast ... processes with summable [exponentially decaying] correlations ... are also called processes with **short memory** or short-range correlations or weak dependence”

How intuitive is the term “memory”?



- The panels on the left show different time windows of the same time series
- The panel above shows how autocorrelation increases with window length, starting from about 0 at the smallest window
- Does the high autocorrelation reflect **memory** or **amnesia**?

Notes on the construction of the previous example

- The time series was constructed by superposition of:
 - A stochastic process with values $\underline{m}_j \sim N(2, 0.5)$ each lasting a period $\underline{\tau}_j$ exponentially distributed with $E[\underline{\tau}_j] = 50$ (red line);
 - White noise $N(0, 0.2)$
- The process of our example is completely **stationary**
- Thus, the autocorrelation function is **independent of time location** and the autocorrelogram is fully meaningful
- However, an estimate thereof from a few terms of a time series is inaccurate
- Nothing in the process reminds us of any type of **memory**
- The high autocorrelation is a result of the **changing** local average

Slow decaying autocorrelation does not reflect long-term memory but long-term change

- The first who pointed this out was **Vit Klemes** (1974)
 - He wrote “It is shown that the **Hurst phenomenon** is not necessarily an indicator of infinite memory of a process”
 - He compared the memory-based interpretation to the Ptolemaic planetary model, which worked well but hampered progress in astronomy for centuries
 - Unfortunately, he used the term “nonstationarity” for change, while in fact his final explanatory model was stationary (and he noted it)
- The Hurst phenomenon, named after **Hurst** (1951) who studied several geophysical time series, essentially reflects long-term changes, even though Hurst formulated it in a slightly different manner:
 - “Although in random events groups of high or low values do occur, their tendency to occur in natural events is greater”
- Earlier than Hurst, **Kolmogorov** (1940) had proposed a mathematical model that can describe stochastically this natural behaviour
- Here the name **Hurst-Kolmogorov (HK) dynamics** is used instead of the term “long-term memory” and the mathematical model is termed the **HK process** (from Koutsoyiannis and Cohn, 2008)



2. What is climate?

A typical definition and some remarks

- “**Climate** [is] the **long-term average of conditions** in the atmosphere, ocean, and ice sheets and sea ice described by **statistics, such as means and extremes**” (U.S. Global Change Research Program, 2009)
- Remark 1: The definition may have missed the importance of the occurrence, circulation, and phase transition of water in the land and the atmosphere, as well the fact that water is the regulator of the entire climate system; hence, it may misrepresent the links of **Climate** with **Hydrology**
- Remark 2: To study “**statistics, such as means and extremes**” one may need some knowledge of **Statistics**
- Remark 3: To study “**long-term average conditions**” one may need to shape multi-scale representations of processes and, hence, one may need some more **advanced** knowledge of **Statistics**
- Remark 4: Inappropriate representations of hydroclimatic processes, based on **Classical Statistics** (assuming independence in time), abound; these have severe consequences in the perception of climate, as well as in hypothesis testing, estimation and prediction

How do we expect climate to look like?



- Both time series are synthetic and have the same marginal properties:
 - Size $n = 2^{16} = 65536$
 - Mean $\mu = 5$
 - Standard deviation $\sigma = 1$
 - Normal distribution
- In both time series “climatic” averages for scales 2^5 and 2^{10} are also given in addition to the “annual” plot



HKC is a non-static, all-scales-changing climate

- Upper time series:
 - Pure randomness (no dependence in time)
 - Change occurs only at scale 1
 - Climatic (i.e. long-term) averages are flat
- Lower time series:
 - HK dependence with Hurst coefficient $H = 0.99$ (see below)
 - Change occurs at all scales
- Real world evidence (see below) supports HKC rather than SC

If climate is ever changing, why coin a term “climate change”?



3. What is the HK process, how it describes the multi-scale properties of a process, and which are its consequences in Statistics?

The Hurst-Kolmogorov (HK) process and its multi-scale stochastic properties

- The HK process does not provide a “perfect” and “detailed” mathematical tool for geophysical processes
- Rather it is the **most parsimonious** and **simplest** alternative to the classical, independence-based, statistical model:
 - It can represent a non-static climate using a single parameter, the Hurst coefficient H , additional to those of classical statistics
 - It produces very simple expressions for all scales
- Also, it is ideal for the perception and intuition development of a non-static climate

Properties of the HK process	At an arbitrary observation scale $k = 1$ (e.g. annual) Process: \underline{x}_i	At any scale k Process: $\underline{x}_i^{(k)} := \frac{1}{k} \sum_{l=(i-1)k+1}^{ik} \underline{x}_l$
Standard deviation	$\sigma \equiv \sigma^{(1)}$	$\sigma^{(k)} = k^{H-1} \sigma$ (H is the Hurst coefficient; $0.5 < H < 1$)
Autocorrelation function (for lag j)	$\rho_j \equiv \rho_j^{(1)} = \rho_j^{(k)} \approx H(2H-1) j ^{2H-2}$	
Power spectrum (for frequency ω)	$S(\omega) \equiv S^{(1)}(\omega) \approx 4(1-H) \sigma^2 (2\omega)^{1-2H}$	$S^{(k)}(\omega) \approx 4(1-H) \sigma^2 k^{2H-2} (2\omega)^{1-2H}$

Can serve as a definition of the HK process

All properties are power laws of scale k , lag j , frequency ω

Impacts on statistical estimation: Hurst-Kolmogorov Statistics (HKS) vs. Classical Statistics (CS)

True values →	Mean, μ	Standard deviation, σ	Autocorrelation ρ_l for lag l
Standard estimator	$\bar{x} := \frac{1}{n} \sum_{i=1}^n x_i$	$\underline{s} := \sqrt{\frac{1}{n-1}} \sqrt{\sum_{i=1}^n (x_i - \bar{x})^2}$	$\underline{r}_l := \frac{1}{(n-1)\underline{s}^2} \sum_{i=1}^{n-l} (x_i - \bar{x})(x_{i+l} - \bar{x})$
Relative bias of estimation, CS	0	≈ 0	≈ 0
Relative bias of estimation, HKS	0	$\approx \sqrt{1 - \frac{1}{n'}} / \sqrt{1 - \frac{1}{n}} - 1 \approx -\frac{1}{2n'}$	$\approx -\frac{1/\rho_l - 1}{n' - 1}$
Standard deviation of estimator, CS	$\frac{\sigma}{\sqrt{n}}$	$\approx \frac{\sigma}{\sqrt{2(n-1)}}$	
Standard deviation of estimator, HKS	$\frac{\sigma}{\sqrt{n'}}$	$\approx \frac{\sigma \sqrt{(0.1n + 0.8)^{\lambda(H)} (1 - n^{2H-2})}}{\sqrt{2(n-1)}}$ where $\lambda(H) := 0.088 (4H^2 - 1)^2$	

Note: $n' := \sigma^2 / \text{Var}[\bar{x}] = n^{2-2H}$ is the “equivalent” or “effective” sample size: a sample with size n' in CS results in the same uncertainty of the mean as a sample with size n in HKS.


See Koutsoyiannis (2003) for derivations and for more accurate expressions.

The variance of the estimator of mean

Definition

$$n' := \sigma^2 / \text{Var}[\bar{x}] = n^{2-2H}$$


Classical statistics (CS)
is here ($H = 0.5$)



An example for
 $H = 0.9$ and $n = 100$


- The equivalent (in a CS sense) sample size $n' = 100^{0.2} = 2.5!$
- The uncertainty in the estimation of mean is $n/n' = 40$ times greater than in CS
- To make the uncertainty equal to that in CS for sample size 100 (i.e. to make $n' = 100$) we need a real sample size $n = 100^{1/0.2} = 10^{10}!$

The bias of the classical estimator of standard deviation



The bias of the classical estimator of autocorrelation

An example for $H = 0.9$ and $n = 100$: Huge bias at all lags



Alternative tools to explore the HK behaviour in time series

- Autocorrelogram?
 - Problematic, because of too high bias
- Periodogram?
 - Very problematic, because it is difficult to obtain analytical expressions of bias and uncertainty, and it may be too rough/scattered (unless the sample size is very high)
- Rescaled range? (the original Hurst's method)
 - Most inappropriate (too much dispersion, high biases, too uncertain estimates, lack of analytical expressions of statistics; see Koutsoyiannis 2002, 2003)
- Residuals of variance (Taqqu *et al.*, 1995) also termed detrended fluctuation analysis (DFA; Peng *et al.*, 1994)
 - Problematic, as it treats biased and uncertain quantities as if they were unbiased and certain
- Climacogram? (term coined in Koutsoyiannis, 2010)
 - Better, provided that we are aware of (and adapt for) biases and uncertainties, for which there exist **analytical expressions**

A typical example of misuse in studies detecting HK behaviour

- Several studies based on exploration of the periodogram, the rescaled range, or the DFA, detect a Hurst coefficient $H > 1$
- This is mathematically absurd, because in a (stationary) HK process (also known as fractional Gaussian noise), $0 \leq H \leq 1$
- Even in a **cumulative** (and thus nonstationary) HK process (also known as fractional Brownian noise), again $0 \leq H \leq 1$ (the cumulative nonstationary process is characterized by the same H as its corresponding stationary process)
- The reported $H > 1$ values are obtained because of inappropriate estimators (based e.g. on slopes of graphs of the explored quantities); their typical interpretation should be that H must be close to 1

See additional discussion in Tyrallis and Koutsoyiannis (2010), which includes:


- Two versions of climacogram-based estimation of the process parameters
- Streamlining of a precise Maximum Likelihood method for parameter estimation
- Comparisons of the three proposed methods as well as intercomparisons with other popular methods

Change and scale: The climacogram

- This is simply a (logarithmic) plot of the standard deviation $\sigma^{(k)}$ at scale k vs. the scale k
- $\sigma^{(k)}$ can be calculated directly from the time averaged process $\underline{x}_i^{(k)}$
- $\sigma^{(k)}$ is a simple transformation of the autocorrelogram ρ_j (where j is lag), i.e.:

$$\sigma^{(k)} = \frac{\sigma}{\sqrt{k}} \sqrt{\alpha_k}, \quad \alpha_k = 1 + 2 \sum_{j=1}^{k-1} \left(1 - \frac{j}{k}\right) \rho_j \quad \leftrightarrow \quad \rho_j = \frac{j+1}{2} \alpha_{j+1} - j \alpha_j + \frac{j-1}{2} \alpha_{j-1}$$

- In classical statistics, $\sigma^{(k)} = \sigma/\sqrt{k}$, which indicates a straight line with slope -0.5 in the logarithmic plot
- In general, the asymptotic slope (for high k) in the logarithmic plot determines the Hurst coefficient:
 $H = 1 + \text{slope}$
- Slopes milder than -0.5, that is, H values in the interval (0.5, 1), indicate long-range dependence



An important note on the use of the climacogram


- In the HK process, the standard deviation $\sigma^{(k)}$ at scale k is related to that at scale 1, i.e. σ , by

$$\sigma^{(k)} := \frac{\sigma}{k^{1-H}}$$

- However, due to bias, the sample standard deviations are not directly comparable to $\sigma^{(k)}$ but to the quantities

$$\sigma_n^{r(k)} := \frac{\sigma}{k^{1-H}} \sqrt{\frac{1 - (k/n)^{2-2H}}{1 - (k/n)}}$$

- Explanation: if $(\underline{s}^{(k)})^2$ is the standard estimator of the variance at scale k , then it can be shown that $E[(\underline{s}^{(k)})^2] = (\sigma_n^{r(k)})^2$
- The example above was constructed assuming a sample with size $n = 100$ from an HK process with $H = 0.9$
- Notice that the slope of the curve “adapted for bias” at scale 10 is -0.21 instead of -0.1 (that of the theoretical climacogram)





4. Does empirical evidence support the hypothesis of HKC?

Example 1: Annual minimum water levels of the Nile





Roda
Nilometer

- The longest available instrumental hydroclimatic data set (813 years)
- Hurst coefficient $H = 0.84$
- The same H is estimated from the simultaneous record of maximum water levels and from the modern record (131 years) of the Nile flows at Aswan



Example 2: The lower tropospheric temperature



Suggests an HK behaviour with a very high Hurst coefficient:
 $H \approx 0.99$



Data: 1979-2009, from
http://vortex.nsstc.uah.edu/public/msu/t2lt/tltghmam_5.2

Example 3: The Monthly Atlantic Multidecadal Oscillation (AMO) Index



Suggests an HK behaviour with a very high Hurst coefficient: $H \approx 0.99$



Data: 1856-2009, from NOAA,
<http://www.esrl.noaa.gov/psd/data/timeseries/AMO/>

Example 4: The annual rainfall in Maatsuyker Island (Australia)

Suggests an HK behaviour with a very high Hurst coefficient:
 $H \approx 0.99$





Maatsuyker Island Lighthouse (Australia),
coordinates: -43.65N, 146.27E, 147 m,
WMO station code: 94962
Data: 1892-2004, available from
<http://climexp.knmi.nl/data/pa94962.dat>

Example 5: The mean annual temperature at Svalbard (Norway)

Suggests an HK behaviour with a high Hurst coefficient:

$$H \approx 0.95$$





Svalbard Luft (Norway),
coordinates: 78.25N, 15.47E, 29 m,
WMO station code: 1008 SVALBARD_LUFT
Data: 1912-2009, available from
<http://www.unaami.noaa.gov/analyses/sat/sat01008.d> and
<http://climexp.knmi.nl/data/ta1008.dat>

Example 6: Iowa fine resolution rainfall



Seven storm events of high temporal resolution, recorded by the Hydrometeorology Laboratory at the Iowa University (Georgakakos *et al.*, 1994)

Suggests an HK behaviour with a very high Hurst coefficient: $H \approx 0.99$



Example 7: The Greenland temperature proxy during the Holocene

Suggests an HK behaviour with a high Hurst coefficient: $H \approx 0.94$



Reconstructed from the GISP2 Ice Core (Alley, 2000, 2004)
Data from:
ftp.ncdc.noaa.gov/pub/data/paleo/icecore/greenland/summit/gisp2/isotopes/gisp2_temp_accum_alley2000.txt


5. Does long-range dependence in the HKC imply lower predictive uncertainty?

Unconditional and conditional entropy in HK

For scale $k = 1$ (say annual), the unconditional uncertainty of a HKC prediction is higher than in RC, but that conditional on the known past is lower

For scale $k = 2$, even the conditional uncertainty of a HKC prediction is higher than in RC


For a climatic scale $k = 30$, the unconditional and conditional standard deviations in HKC are respectively 5 and 3 times higher than in RC



- The graph was constructed using analytical equations by Koutsoyiannis (2005) and Koutsoyiannis *et al.* (2007)
- The following assumptions were made:
 - $H = 0.90$ (known a priori)
 - Sample size $n = 100$
 - Sample standard deviation $s = 1$
 - Normal distribution

Generalized comparison of HKC with RC predictions


- The graph was constructed with same assumptions as the previous one but with varying sample size n and Hurst coefficient H , which is assumed *a priori* known
- In reality, the results are worse because H is estimated from data
- Practical conclusion: except for scale 1 (e.g. annual) the HK behaviour increases predictive uncertainty



The curves show the scale k in which the HKC uncertainty (conditional on known past) becomes greater than in RC

Does switching from stochastic to deterministic description of climate enable long term predictability?

- The Figure on the right shows “**Long-term Earth system projections** for different CO₂ emission and storage scenarios” (from Shaffer, 2010)
 - They refer to several climatic variables including “mean atmosphere warming” and “mean ocean warming”
 - They extend up to **100 000 A.D.** (!)
 - The information provided is supposed to support “geoengineering” options, e.g. the “effectiveness and consequences of carbon dioxide sequestration”
- Question 1: Do deterministic models reproduce the past? (see my answer in Koutsoyiannis *et al.*, 2008, Anagnostopoulos *et al.*, 2010)
- Question 2: Does knowledge of deterministic dynamics enable reliable future predictions? (see my answer in Koutsoyiannis, 2010)



6. Conclusion

Concluding remarks

- Change is Nature's style and occurs at all time scales
- Climate is no exception: it is ever changing
- By definition, the very notion of climate relies on Statistics—in particular, on long-term statistical properties of natural processes
- Classical Statistics is inconsistent with real-world climate; rather, it corresponds to a static climate
- Hurst-Kolmogorov Statistics is the key to perceive a multi-scale changing climate and model the implied uncertainty and risk
- Hurst-Kolmogorov Statistics has marked differences from Classical Statistics, which are often missed; thus, it implies:
 - Dramatically higher uncertainty of statistical parameters of location
 - High negative bias of statistical parameters of dispersion and dependence
 - Dramatically higher predictive uncertainty

“Υβριν χρή σβεννύναι μᾶλλον ἢ πυρκαϊήν (Ἡράκλειτος, ἀπόσπασμα 43)
More than a fire, one needs to extinguish hubris (Heraclitus, fragment 43)

References

- Alley, R. B., The Younger Dryas cold interval as viewed from central Greenland, *Quaternary Science Reviews*, 19, 213-226, 2000
- Alley, R. B., GISP2 Ice Core Temperature and Accumulation Data, IGBP PAGES/World Data Center for Paleoclimatology Data Contribution Series #2004-013, NOAA/NGDC Paleoclimatology Program, Boulder CO, USA, 2004
- Anagnostopoulos, G. G., D. Koutsoyiannis, A. Christofides, A. Efstratiadis, and N. Mamassis, A comparison of local and aggregated climate model outputs with observed data, *Hydrological Sciences Journal*, 2010 (in press)
- Beran, J., *Statistics for Long-Memory Processes*, Vol. 61 of Monographs on Statistics and Applied Probability, Chapman and Hall, New York, 1994
- Georgakakos, K., A. Carsteanu, P. Sturdevant, and J. Cramer, Observation and Analysis of Midwestern Rain Rates, *Journal of Applied Meteorology*, 33(12), 1433-1444, 1994
- Hurst, H. E., Long term storage capacities of reservoirs, *Trans. Am. Soc. Civil Engrs.*, 116, 776–808, 1951
- Klemes, V., The Hurst phenomenon: A puzzle?, *Water Resour. Res.*, 10(4) 675-688, 1974
- Kolmogorov, A. N., Wienersche Spiralen und einige andere interessante Kurven in Hilbertschen Raum, *Dokl. Akad. Nauk URSS*, 26, 115–118, 1940
- Koutsoyiannis, D., The Hurst phenomenon and fractional Gaussian noise made easy, *Hydrological Sciences Journal*, 47 (4), 573–595, 2002
- Koutsoyiannis, D., Climate change, the Hurst phenomenon, and hydrological statistics, *Hydrological Sciences Journal*, 48 (1), 3–24, 2003
- Koutsoyiannis, D., Uncertainty, entropy, scaling and hydrological stochastics, 2, Time dependence of hydrological processes and time scaling, *Hydrological Sciences Journal*, 50 (3), 405–426, 2005
- Koutsoyiannis, D., A random walk on water, *Hydrology and Earth System Sciences*, 14, 585–601, 2010
- Koutsoyiannis, D., and T.A. Cohn, The Hurst phenomenon and climate (solicited), *European Geosciences Union General Assembly 2008, Geophysical Research Abstracts, Vol. 10*, Vienna, 11804, European Geosciences Union, 2008 (<http://www.itia.ntua.gr/en/docinfo/849/>)
- Koutsoyiannis, D., A. Efstratiadis, and K. Georgakakos, Uncertainty assessment of future hydroclimatic predictions: A comparison of probabilistic and scenario-based approaches, *Journal of Hydrometeorology*, 8 (3), 261–281, 2007
- Koutsoyiannis, D., A. Efstratiadis, N. Mamassis, and A. Christofides, On the credibility of climate predictions, *Hydrological Sciences Journal*, 53 (4), 671–684, 2008
- Papoulis, A., *Probability, Random Variables, and Stochastic Processes*, 3rd ed., McGraw-Hill, New York, 1991
- Peng, C. K., S. V. Buldyrev, M. Simons, H. E. Stanley and A. L. Goldberger, Mosaic organization of DNA nucleotides, *Physical Review E*, 49, 1685-1689, 1994
- Shaffer, G., Long-term effectiveness and consequences of carbon dioxide sequestration, *Nature Geoscience*, 3, 464 – 467, 2010
- Taqqu, M., V. Teverovsky and W. Willinger, Estimators for long-range dependence: an empirical study, *Fractals* 3 (4), 785-798, 1995
- Tyralis, H., and D. Koutsoyiannis, Simultaneous estimation of the parameters of the Hurst-Kolmogorov stochastic process, *Stochastic Environmental Research & Risk Assessment*, 2010 (in press)
- U.S. Global Change Research Program, *Climate Literacy, The Essential Principles of Climate Sciences*, 2nd version, 2009 (<http://downloads.climate-science.gov/Literacy/Climate%20Literacy%20Booklet%20Low-Res.pdf>)